

- [4] Arnold, C. A., 1938, 1948: Paleozoic seeds. *Bot. Rev.*, 4(5): 205—234, 14(7): 450—472.  
 [5] Critchfield, W. B., 1970: Shoot growth and heterophylly in *Ginkgo biloba*. *Bot. Gaz.*, 131: 150—162.  
 [6] Fujii, K., 1896: On the different views hitherto proposed regarding the morphology of the flowers of *Ginkgo biloba*. *Bot. Mag. (Tokyo)*, 10: 15—25, 104—110.  
 [7] Sakisaka, M., 1929: On the seedbearing leaves of *Ginkgo*. *Jap. J. Bot.*, 4.  
 [8] Wilson, C. L., 1953: The telome theory. *Bot. Rev.*, 19: 417—437.

## 中国鬼灯檠属新分类群

### NEW TAXA OF THE GENUS *RODGERSIA* FROM CHINA

**Abstract** A new status, *Rodgersia aesculifolia* var. *henricii* (Franch.) C.Y.Wu, is proposed, and two new varieties, i. e. *R. sambucifolia* var. *estrigosa* J.T. Pan, and *R. pinnata* var. *strigosa* J. T. Pan are described in this paper.

**Key words** *Rodgersia*; new taxa

**关键词** 鬼灯檠; 新分类群

#### 滇西鬼灯檠 新等级

***Rodgersia aesculifolia* Batalin var. *henricii* (Franch.) C. Y. Wu, stat. nov. —**  
*Rodgersia henricii* (Franch.) Franch. in *Rev. Hort. Paris* 69: 176. 1897.

**Yunnan** (云南): Weixi (维西), Yezhi (叶枝), on grassy slopes, alt. 3600 m, 1935. 08, C. W. Wang (王启无) 68522.

#### 光腹鬼灯檠 新变种

***Rodgersia sambucifolia* Hemsl. var. *estrigosa* J. T. Pan, var. nov.**

A var. *sambucifolia* differt laminis foliolorum ventre glabris.

**Yunnan** (云南): Lijiang (丽江), on mountain slopes, alt. 2300 m, 1935. 06, C. W. Wang (王启无) 70635 (holotype, PE)

此变种与原变种之区别在于其小叶片腹面无毛。

#### 伏毛鬼灯檠 新变种

***Rodgersia pinnata* Franch. var. *strigosa* J. T. Pan, var. nov.**

A var. *pinnata* recedit lamellis foliolorum ventre strigosis.

**Sichuan** (四川): Meigu (美姑), on slopes, 1959. 08. 21, Exp. Econ. Pl. Sich. (四川经济植物采集队) 1893 (holotype, PE)

本变种不同原变种在于小叶片腹面被糙伏毛。

潘锦堂 (中国科学院西北高原生物研究所, 西宁 810001) Pan Jin-tang  
 (North-western Plateau Institute Biology, Academia Sinica, Xining 810001)